

Triggering collective oscillations by three-flavor effects

Basudeb Dasgupta,¹ Georg G. Raffelt,¹ and Irene Tamborra^{1,2,3}

¹*Max-Planck-Institut für Physik (Werner-Heisenberg-Institut), Föhringer Ring 6, 80805 München, Germany*

²*Dipartimento Interateneo di Fisica “Michelangelo Merlin,” Via Amendola 173, 70126 Bari, Italy*

³*Istituto Nazionale di Fisica Nucleare, Sezione di Bari, Via Orabona 4, 70126 Bari, Italy*

Collective flavor transformations in supernovae, caused by neutrino-neutrino interactions, are essentially a two-flavor phenomenon driven by the atmospheric mass difference and the small mixing angle θ_{13} . In the two-flavor approximation, the initial evolution depends logarithmically on θ_{13} and the system remains trapped in an unstable fixed point for $\theta_{13} = 0$. However, any effect breaking exact ν_μ - ν_τ equivalence triggers the conversion. Such three-flavor perturbations include radiative corrections to weak interactions, small differences between the ν_μ and ν_τ fluxes, or non-standard interactions. Therefore, extremely small values of θ_{13} are in practice equivalent, the fate of the system depending only on the neutrino spectra and their mass ordering.

PACS numbers: 14.60.Pq, 97.60.Bw

I. INTRODUCTION

Dense neutrino ensembles exhibit collective flavor transformations [1–6]. For supernova (SN) neutrinos, these effects imprint intriguing features on the processed spectrum such as spectral splits [7–19]. Collective oscillations originate from a characteristic instability in flavor space of the interacting dense neutrino system that partly swaps its flavor content to minimize its energy. Assuming the primary ν_e and $\bar{\nu}_e$ fluxes exceed those of the other flavors, the dominant effect arises for inverted neutrino mass hierarchy. It is driven by the atmospheric mass difference and the mixing angle θ_{13} that is known to be small and could vanish entirely.

The transformation arising from an instability implies that the processed spectrum is independent of the mixing angle as long as it is small (collective transformations in the presence of matter are suppressed for maximal mixing because one projects on the interaction direction, but this $\cos 2\theta$ effect is irrelevant if $\theta \ll 1$ [20, 22]). In a two-flavor treatment, θ enters only as a trigger to the subsequent evolution, so in the SN context a very small θ shifts logarithmically the onset radius for collective transformations [8, 9]. In numerical studies, choosing θ as small as allowed by the machine precision barely impacts the processed spectrum, although for $\theta = 0$ the system remains stuck in the unstable fixed-point solution defined by the initial conditions.

Such a situation looks unphysical—it does not seem plausible that, at least in principle, one can distinguish between θ being exactly zero and some arbitrarily small but non-zero value. One may speculate, for example, that quantum fluctuations could trigger the transformation even for $\theta = 0$ [8], noting that collective transformations actually preserve flavor lepton number.

The purpose of our paper is to show that in real life we do not need to worry about such subtleties. If θ_{13} is sufficiently small, three-flavor effects [20, 23–27] trigger the instability and the logarithmic θ_{13} dependence saturates at a small but non-zero value.

Why are collective SN neutrino transformations an effective two-flavor phenomenon anyway? In the outer layers the temperature is too low to support thermal μ or τ populations, obviating the possibility to distinguish between ν_μ and ν_τ by charged-current reactions. Ignoring radiative corrections, these flavors are exactly equivalent, allowing us to define new flavors ν_x and ν_y such that effectively $\theta_{23} = 0$. If in addition $\theta_{13} = 0$, one of the new states, say ν_y , becomes equivalent to ν_3 , decoupling entirely from the other flavors: We are left with a two-flavor system consisting of ν_e and ν_x , governed by θ_{12} and the solar mass difference δm^2 .

This system can show collective transformations. However, the solar mass hierarchy is normal, suppressing the dominant transformation effect if the primary fluxes show the usual excess of ν_e and $\bar{\nu}_e$. Moreover, the collective oscillation region is at larger radii because the solar mass difference is small, so multiple-split effects are more easily suppressed by adiabaticity violation [12, 13]. However, collective oscillations driven by the solar mass difference do modify the spectra in some scenarios [21].

Once we allow for a small but non-vanishing θ_{13} , collective $\nu_e \leftrightarrow \nu_y$ transformations become possible that are driven by the atmospheric mass difference Δm^2 and occur in the usual region of large neutrino flux densities. Our main point is that for $\theta_{13} = 0$ these transformations are triggered by small perturbations of the exact ν_μ - ν_τ equivalence because ν_e and ν_x then no longer form an exact two-flavor system. Such perturbations include radiative corrections to the ν_μ and ν_τ matter effect [28, 29], or small ν_μ - ν_τ flux differences. The latter can be caused by the presence of muons in deeper layers of the SN core and by radiative corrections to the interaction rates, modifying the relative ν_μ and ν_τ opacities. Non-standard interactions can also break the ν_μ - ν_τ symmetry [27], a possibility that we will not pursue here.

We begin in Sec. II with a brief discussion of the equations of motion. In Sec. III we prove that for $\theta_{13} = 0$ and for exact ν_μ - ν_τ equivalence, collective oscillations driven by the atmospheric mass difference are not possible, justifying the usual two-flavor treat-

ment. In Sec. IV we study concrete departures from ν_μ - ν_τ equivalence in the limit $\theta_{13} = 0$ and show that collective transformations are triggered by these effects. In schematic models we compare them with an equivalent θ_{13} that would trigger collective transformations at the same onset radius. In Sec. V we consider a realistic SN and study the competition between a small θ_{13} and a small ν_μ - ν_τ flux difference. We conclude with a brief summary in Sec. VI.

II. EQUATIONS OF MOTION

A. Matrix form

For our conceptual discussion it is sufficient to consider the simplest three-flavor system showing collective transformations. We take the neutrino ensemble to be homogeneous and isotropic, study its time evolution, and describe mixed neutrinos by matrices of densities ϱ_E for each energy mode E . We use an overbar to represent the corresponding quantities for antineutrinos. Diagonal entries are the usual occupation numbers, whereas off-diagonal entries encode phase information. The equations of motion (EoM) are

$$i \dot{\varrho}_E = [\mathbf{H}_E, \varrho_E] \quad \text{and} \quad i \dot{\bar{\varrho}}_E = [\bar{\mathbf{H}}_E, \bar{\varrho}_E]. \quad (1)$$

The Hamiltonian 3×3 matrix is made up of the vacuum, matter, and neutrino-neutrino terms

$$\mathbf{H}_E = \mathbf{H}_E^{\text{vac}} + \mathbf{H}^\lambda + \mathbf{H}^{\nu\nu}. \quad (2)$$

Here $\mathbf{H}_E^{\text{vac}} = \mathbf{U} \mathbf{M}^2 \mathbf{U}^\dagger / 2E$, with $\mathbf{U} = \mathbf{R}_{23} \mathbf{R}_{13} \mathbf{R}_{12}$ the neutrino mixing matrix and $\mathbf{M} = \text{diag}(m_1, m_2, m_3)$ the mass matrix. We use the standard notation of $\mathbf{R}_{ij}$ as the rotation matrix between the i and j mass eigenstates, with argument θ_{ij} . For antineutrinos, the vacuum Hamiltonian picks up a relative minus sign ($\bar{\mathbf{H}}_E^{\text{vac}} = -\mathbf{H}_E^{\text{vac}}$), whereas all other pieces remain identical. For oscillation studies, we may neglect terms proportional to the identity and may write in the mass basis

$$\mathbf{H}_E^{\text{vac}} = \frac{1}{2E} \text{diag} \left(-\frac{\delta m^2}{2}, +\frac{\delta m^2}{2}, \Delta m^2 \right). \quad (3)$$

The solar mass-squared difference $\delta m^2 > 0$, whereas the atmospheric one $\Delta m^2 < 0$ for inverted mass hierarchy (IH) and $\Delta m^2 > 0$ for normal hierarchy (NH).

The matter term, due to neutrino interactions with the charged leptons, is in the flavor basis

$$\mathbf{H}^\lambda = \sqrt{2} G_F \text{diag}(N_e, N_\mu, N_\tau) + \mathcal{O}(G_F^2), \quad (4)$$

where N_e is the net electron density (electrons minus positrons) and similarly for the other leptons. The second-order term is due to radiative corrections and can be non-negligible at high densities [28]. The contribution associated to ν - ν interactions is

$$\mathbf{H}^{\nu\nu} = \sqrt{2} G_F \int dE (\varrho_E - \bar{\varrho}_E) + \mathcal{O}(G_F^2). \quad (5)$$

Multi-angle effects are ignored in our isotropic system. Radiative corrections can be important in dense neutrino gases at the second order [29].

B. New interaction basis e - x - y

Since we are concerned with a system where the ν_μ and ν_τ flavors are exactly or approximately equivalent, it is more useful to introduce new flavors x and y that simplify the mixing matrix [24]

$$\begin{pmatrix} \nu_e \\ \nu_x \\ \nu_y \end{pmatrix} = \mathbf{R}_{23}^\dagger \begin{pmatrix} \nu_e \\ \nu_\mu \\ \nu_\tau \end{pmatrix}. \quad (6)$$

Here $\mathbf{R}_{23}^\dagger$ “unmixes” ν_μ and ν_τ with the angle θ_{23} . For $\theta_{13} = 0$, ν_y is the mass eigenstate ν_3 . Henceforth the interaction basis is understood to be the e - x - y -basis.

This basis is useful because it explicitly removes θ_{23} from the formalism, if the Hamiltonian and the initial conditions do not distinguish ν_μ and ν_τ . Naturally, the evolution of ν_e and $\bar{\nu}_e$ is independent of θ_{23} in this approximation.

C. Expansion in Gell-Mann matrices

The commutator structure of the equations of motion ensures that the trace of ϱ_E is conserved, so we may re-define them to be traceless by subtracting a term proportional to the identity matrix $\mathbf{I}$. The traceless part can be expanded in Gell-Mann matrices Λ_i with the expansion coefficients forming an 8-vector $\mathbf{X}$. Thus one can project any matrix $\mathbf{X}$ as

$$\mathbf{X} = \text{tr}(\mathbf{X}) \left(\frac{\mathbf{I}}{3} + \mathbf{X} \cdot \frac{\boldsymbol{\Lambda}}{2} \right), \quad (7)$$

where $\boldsymbol{\Lambda}$ is an 8-vector of Λ matrices. We normalize as $|\mathbf{X}| = 2/\sqrt{3}$, corresponding to the conventions $\Lambda_i = \Lambda_i^\dagger$, $\text{tr}(\Lambda_i) = 0$, $\text{tr}(\Lambda_i \Lambda_j) = 2\delta_{ij}$, and $[\Lambda_i, \Lambda_j] = 2if_{ijk}\Lambda_k$. Here f_{ijk} are the $\text{SU}(3)$ structure constants, where

$$\begin{aligned} f_{123} &= 1 \\ f_{147} &= f_{246} = f_{257} = f_{345} = -f_{156} = -f_{367} = 1/2 \\ f_{458} &= f_{678} = \sqrt{3}/2 \end{aligned} \quad (8)$$

are the non-vanishing values.

The neutrino matrices of density can now be decomposed, as in Eq. (7), in terms of an 8-dimensional polarization vector

$$\varrho_E = n_E \left(\frac{\mathbf{I}}{3} + \mathbf{P}_E \cdot \frac{\boldsymbol{\Lambda}}{2} \right). \quad (9)$$

n_E is the total neutrino density per unit energy interval. Analogous expressions pertain to antineutrinos.

The different parts of the Hamiltonian can also be similarly decomposed. The vacuum Hamiltonian is

$$H_E^{\text{vac}} = \omega_E \left(\frac{\mathbf{I}}{3} + \mathbf{B} \cdot \frac{\mathbf{\Lambda}}{2} \right), \quad (10)$$

where $\omega_E = \Delta m^2/(2E)$. The “magnetic field” is

$$\mathbf{B} = \begin{pmatrix} 0 \\ 0 \\ s_{13}^2 \\ S_{13} \\ 0 \\ 0 \\ 0 \\ -\frac{1}{2\sqrt{3}}(1+3C_{13}) \end{pmatrix} + \epsilon_\omega \begin{pmatrix} c_{13}S_{12} \\ 0 \\ -\frac{1}{4}C_{12}(3+C_{13}) \\ \frac{1}{2}C_{12}S_{13} \\ 0 \\ -s_{13}S_{12} \\ 0 \\ \frac{\sqrt{3}}{2}C_{12}s_{13}^2 \end{pmatrix}, \quad (11)$$

with $\epsilon_\omega = \delta m^2/\Delta m^2$. Moreover, we use $s_{ij} = \sin \theta_{ij}$, $c_{ij} = \cos \theta_{ij}$, $S_{ij} = \sin 2\theta_{ij}$, and $C_{ij} = \cos 2\theta_{ij}$.

Ignoring a term proportional to identity, the matter term can be written as

$$H^\lambda = \lambda \text{diag}(1, 0, \epsilon_\lambda) = \lambda \left(\frac{\mathbf{I}}{3} + \mathbf{L} \cdot \frac{\mathbf{\Lambda}}{2} \right). \quad (12)$$

Here $\lambda = \sqrt{2}G_F N_e$ is the effective MSW potential. For later reference we have included $\epsilon_\lambda \ll 1$, encoding radiative corrections or small ν_μ - ν_τ flux differences. The leptonic “magnetic field” is

$$\mathbf{L} = \begin{pmatrix} 0 \\ 0 \\ 1 \\ 0 \\ 0 \\ 0 \\ 0 \\ \frac{1}{\sqrt{3}} \end{pmatrix} + \epsilon_\lambda \begin{pmatrix} 0 \\ 0 \\ -s_{23}^2 \\ 0 \\ 0 \\ -S_{23} \\ 0 \\ -\frac{1}{2\sqrt{3}}(1+3C_{23}) \end{pmatrix}. \quad (13)$$

The ν - ν interaction term finally is

$$H^{\nu\nu} = \mu \left(\frac{\mathbf{I}}{3} + \mathbf{D} \cdot \frac{\mathbf{\Lambda}}{2} \right), \quad (14)$$

where the effective neutrino-neutrino interaction energy is $\mu = \sqrt{2}G_F(N + \bar{N})$. Here $N = N_{\nu_e} + N_{\nu_\mu} + N_{\nu_\tau}$ is the overall neutrino density, and $\bar{N}$ for antineutrinos. The collective vector $\mathbf{D}$ is explicitly

$$\mathbf{D} = \int dE \frac{n_E \mathbf{P}_E - \bar{n}_E \bar{\mathbf{P}}_E}{N + \bar{N}}. \quad (15)$$

The EoM are then

$$\dot{\mathbf{P}}_E = (+\omega_E \mathbf{B} + \lambda \mathbf{L} + \mu \mathbf{D}) \times \mathbf{P}_E, \quad (16)$$

$$\dot{\bar{\mathbf{P}}}_E = (-\omega_E \mathbf{B} + \lambda \mathbf{L} + \mu \mathbf{D}) \times \bar{\mathbf{P}}_E. \quad (17)$$

The 8-dimensional vector product is defined as $(\mathbf{a} \times \mathbf{b})_i = f_{ijk} a_j b_k$. In this form, the problem resembles a set of polarization vectors $\mathbf{P}_E$ precessing under the influence of the combined magnetic fields $\mathbf{B}$, $\mathbf{L}$, and the mean field $\mathbf{D}$ due to all polarization vectors.

III. EXACT ν_μ - ν_τ EQUIVALENCE

In the approximation that nothing distinguishes between the ν_μ and ν_τ flavor, 2–3 mixing is physically irrelevant and we expect that oscillations reduce to a two-flavor problem. In fact, for $\theta_{13} = 0$, no collective effects driven by the atmospheric mass difference occur.

To prove this point we study a simplified system consisting of two Bloch vectors, representing equal numbers of neutrinos and antineutrinos, with the single vacuum oscillation frequency ω . For $\theta_{13} = 0$ the magnetic field simplifies to

$$\mathbf{B} = \frac{2}{\sqrt{3}} \mathbf{e}_8 + \epsilon_\omega (S_{12} \mathbf{e}_1 - C_{12} \mathbf{e}_3), \quad (18)$$

where $\mathbf{e}_i$ are unit vectors in the 8-dimensional flavor space. Assuming exact ν_μ - ν_τ equivalence implies that $\epsilon_\lambda = 0$, and therefore

$$\mathbf{L} = \mathbf{e}_3 + \frac{1}{\sqrt{3}} \mathbf{e}_8. \quad (19)$$

Likewise, if the initial ν_μ and ν_τ densities are equal, the initial polarization vectors $\mathbf{P} = \bar{\mathbf{P}}$ are proportional to the same linear combination of $\mathbf{e}_3$ and $\mathbf{e}_8$.

The static vectors $\mathbf{B}$ and $\mathbf{L}$ have components in the 1, 3, and 8 directions, whereas the only dynamical component of $\mathbf{H}$, the self-term $\mathbf{D}$, develops an $\mathbf{e}_2$ component. The EoM of the $\mathbf{D}$ vector derives from the difference of Eqs. (16) and (17)

$$\dot{\mathbf{D}} = -\epsilon_\omega \omega [S_{12}(P_3 + \bar{P}_3) + C_{12}(P_1 + \bar{P}_1)] \mathbf{e}_2. \quad (20)$$

In other words, the vector $\mathbf{H} = \omega \mathbf{B} + \lambda \mathbf{L} + \mu \mathbf{D}$ has only components in the 1, 2, 3 and 8 direction and thus can not mix ν_e and ν_y .

The same conclusion is reached if we consider the EoM in matrix form. The part consisting of the e and x flavor and the y flavor form separate block matrices both for the Hamiltonian matrix and the matrices of densities. In the e - x - y basis and with $\theta_{13} = 0$, the third mass eigenstate ν_3 is not admixed to the ν_e and ν_x flavors.

IV. BROKEN ν_μ - ν_τ EQUIVALENCE

Even in the absence of thermal μ or τ populations the exact ν_μ - ν_τ equivalence is broken by several sub-leading effects that distinguish between these flavors. In this case, the ν_3 flavor does not fully decouple from the ν_e - ν_x system and collective transitions driven by the atmospheric mass difference are inevitably triggered.

The first is provided by radiative corrections to the neutrino matter effect where charged leptons appear in the loop. Even in the absence of ordinary matter, similar radiative corrections arise for neutrino-neutrino interactions, although the detailed structure of the EoM becomes more complicated [29]. Since collective effects

require a large density of neutrinos, radiative corrections and thus the breaking of ν_μ - ν_τ equivalence are unavoidable. Finally we note that differences in the initial ν_μ and ν_τ fluxes also provides the required instability.

A. Radiative corrections to ν_τ matter effect

The presence of matter (i.e. $\lambda \neq 0$) has a similar effect as decreasing the effective mixing angle, although in detail the dynamics is more complicated. In a frame rotating around $\mathbf{L}$ there is a fast-rotating transverse B -field that disturbs the system and triggers the evolution [8]. However, if matter effects distinguish ν_μ and ν_τ , they can play a more important role in triggering collective oscillations, particularly for a small mixing angle. The largest correction is for ν_τ where a background of ordinary matter with baryon density N_B has the same refractive effect on ν_τ and $\bar{\nu}_\tau$ that would be provided by a density of real τ leptons ($N_\tau^{\text{eff}} = 2.6 \times 10^{-5} N_B$) [28]. This subleading correction is parametrized as ϵ_λ to the usual matter effect λ in Eq. (13).

Off-diagonal terms in the Hamiltonian generated by ϵ_λ will mix ν_e and ν_y . In the limit $\theta_{13} \rightarrow 0$, when ν_3 would otherwise have decoupled, these terms play a role similar to θ_{13} , and recouple ν_3 to ν_e . We can estimate the effective θ_{13} generated by these sub-leading matter effects by diagonalizing the Hamiltonian instantaneously in matter as

$$H_E^{\text{vac}} + H^\lambda = \frac{\tilde{U}\tilde{M}^2\tilde{U}^\dagger}{2E}, \quad (21)$$

where the $\tilde{M}$ and $\tilde{U}$ denote the mass and mixing matrix in matter. Using the standard parametrization, Eq. (21) can be solved for the parameters of $\tilde{U}$ and $\tilde{M}$. We find that the matter induced 1–3 mixing is

$$\tan 2\tilde{\theta}_{13} = \frac{2\epsilon_\lambda\epsilon_\omega\lambda S_{12}S_{23}}{2\omega + 2\epsilon_\omega\omega C_{12} + (\epsilon_\lambda - 2)\lambda + \epsilon_\lambda\lambda C_{23}}, \quad (22)$$

where we ignore terms beyond the leading order in ϵ_λ . This equation should be interpreted as providing the critical value of θ_{13} such that if $\theta_{13} \lesssim \tilde{\theta}_{13}$, the role of ν_μ - ν_τ equivalence breaking is more important than θ_{13} itself, and the two-flavor approximation is not valid anymore.

To demonstrate this, we consider a toy model with one Bloch vector for neutrinos $\mathbf{P}$ and one for antineutrinos $\bar{\mathbf{P}}$ with equal length. In the two-flavor case this would be the simple flavor pendulum without intrinsic angular momentum. A nonvanishing mixing angle triggers an exponential growth of the misalignment between the force direction and the initial orientation. The time (or distance) after which an $\mathcal{O}(1)$ deviation from the initial orientation is achieved grows logarithmically with decreasing mixing angle. We define the radius at which there is a change of 1% in the ν_e flavor content as the onset radius. In Fig. 1, we show the onset radius for this system, and how it depends on θ_{13} . We use




FIG. 1: Onset radius as a function of θ_{13} for the toy model described in the text. We assume a small difference between ν_μ and ν_τ refraction parameterized by $\epsilon_\lambda = 10^{-5}$.

$\lambda = 100 \text{ km}^{-1}$, $\epsilon_\lambda = 5 \times 10^{-5}$, $\mu = 10 \text{ km}^{-1}$, $\omega = 1 \text{ km}^{-1}$, and the mixing angles $\theta_{12} = 0.6$, $\theta_{23} = \pi/4$. Using Eq. (22) for the chosen parameters, we expect that the matter induced mixing becomes important at $\theta_{13} \approx 10^{-7}$. This is in good agreement with what we find. The logarithmic increase of the onset radius stops below this critical mixing angle.

B. Different primary ν_μ and ν_τ fluxes

Another way to break the exact ν_μ - ν_τ equivalence is through an initial flux difference. Although this effect is inevitable it has not been studied in detail. Deep in a SN core, the temperature is large enough to support a thermal muon population, slightly modifying the primary fluxes. Moreover, the same radiative effects that create a refractive difference between ν_μ and ν_τ also modify the scattering rates and the two flavors will have slightly different opacities and therefore different thermally driven fluxes. Obviously the discrete nature of particle emission and thermal fluctuations of the regions emitting the neutrinos would necessarily make the two spectra different.

As a toy example we again assume two equal Bloch vectors $\mathbf{P}$ and $\bar{\mathbf{P}}$. The difference between the initial densities of ν_μ and ν_τ is parameterized as

$$\epsilon_N = \frac{N_{\nu_\mu} - N_{\nu_\tau}}{N_{\nu_e}}, \quad (23)$$

and the same for antineutrinos. Ignoring the matter effect, we have explicitly

$$\dot{\mathbf{D}} = \omega \mathbf{B} \times (\mathbf{P} + \bar{\mathbf{P}}), \quad (24)$$

which dynamically generates components of $\mathbf{D}$ along $\mathbf{e}_5$ and $\mathbf{e}_7$ even though initially $\mathbf{H}$ has only components in the 1, 2, 3, and 8 directions. Therefore $\mathbf{H}$ develops a component along $\mathbf{e}_5$, leading to a mixing of ν_e and ν_y .




FIG. 2: Same as Fig. 1, now for equal ν_μ and ν_τ refraction, but for different fluxes as indicated by the values of ϵ_N .

It is not straightforward to define an effective mixing angle in this case. The effect of the different fluxes for ν_μ and ν_τ is to provide terms proportional to $\epsilon_N \mu$ to the $\nu_x - \nu_y$ block in the Hamiltonian. These terms are themselves dynamical (time-dependent), and are communicated to the $\nu_e - \nu_y$ block by the mixing between ν_e and ν_x . The effective mixing angle can be thought as being the initial misalignment of $\mathbf{P}$ from the Hamiltonian which is approximately proportional to $\epsilon_N / (\omega + \lambda)$. As this is a three-flavor effect, it must vanish when $\epsilon_\omega \rightarrow 0$. We therefore expect

$$\tilde{\theta}_{13} \sim \frac{\epsilon_N \epsilon_\omega \omega}{\omega + \lambda}. \quad (25)$$

The logarithmic increase of r_{onset} with decreasing θ_{13} saturates at θ_{13} approximately equal to the effective mixing $\tilde{\theta}_{13}$ induced by unequal $\nu_\mu - \nu_\tau$ fluxes.

In Fig. 2, we plot r_{onset} for this system as a function of θ_{13} for different values of ϵ_N , illustrating this effect. We use the frequencies $(\omega, \mu, \lambda) = (1, 10, 100) \text{ km}^{-1}$, and the mixing angles $\sin^2 \theta_{12} = 0.314$, and $\sin^2 \theta_{23} = 0.5$. Using Eq. (25) for the chosen parameters, we expect the flux-asymmetry induced mixing to become important at $\theta_{13} \sim \epsilon_N / (3 \times 10^{-3})$. This is in good agreement with what we find. The logarithmic increase of the onset radius stops below the estimated value of the mixing angle.

V. REALISTIC SUPERNOVA

We finally consider a more realistic SN example in a single-angle treatment. The neutrinos are assumed to be emitted isotropically from the neutrinosphere at $R_\nu = 10 \text{ km}$. We assume equal luminosities for all neutrino flavors, given by

$$L = 1.2 \times 10^{52} \text{ erg/s}, \quad (26)$$

and thermal spectra with average energies $\langle E_{\nu_e} \rangle = 10$, $\langle E_{\bar{\nu}_e} \rangle = 15$, and $\langle E_{\bar{\nu}_{\mu,\tau}} \rangle = 20 \text{ MeV}$. The electron density

of the matter is the same as in [30] at $t = 1 \text{ s}$ after the bounce. For the neutrino mixing parameters we use

$$\begin{aligned} \Delta m^2 &= 2 \times 10^{-3} \text{ eV}^2, \\ \delta m^2 &= 8 \times 10^{-5} \text{ eV}^2, \\ \sin^2 \theta_{12} &= 0.31, \\ \sin^2 \theta_{23} &= 0.50. \end{aligned} \quad (27)$$

With these assumptions, we have calculated the onset radius for collective transformations as a function of θ_{13} , assuming a flux difference $\epsilon_N = 10^{-5}$ and ignoring radiative corrections to the matter effect. Our results are shown in Fig. 3. For $\theta_{13} \lesssim 10^{-3}$, the onset radius is not sensitive to θ_{13} , as expected from Eq. (25).




FIG. 3: Onset radius of collective oscillations for a realistic SN example, assuming $\epsilon_N = 10^{-5}$.

In a realistic SN, even when the $\nu_\mu - \nu_\tau$ equivalence is perfect, the onset radius depends only very weakly on θ_{13} . The unavoidable breaking of this symmetry by radiative corrections and the presence of charged muons in the deep SN core almost completely removes the θ_{13} dependence in the three-flavor context. Of course, MSW transitions caused by the ordinary matter effect depend on θ_{13} in the usual way.

VI. CONCLUSIONS

Collective oscillations are an instability-driven phenomenon. The system transits from its initial unstable configuration to a stable one, triggered by the influence of a disturbance. Usually one thinks of this disturbance as being provided by the small offset between the relevant flavor and the propagation eigenstates, encoded into the mixing angle θ_{13} . When this mixing angle is exactly vanishing, one would naively think that the oscillations do not take place.

However, one should recognize that a system sitting on an unstable fixed point is bound to be disturbed, unless there are symmetries that forbid all perturbations capable of providing an initial disturbance. In the neutrino

oscillation context, this symmetry happens to be the μ - τ symmetry - which is explicitly broken. Consequently, collective oscillations are inevitable. This means that collective oscillations take place as usual even at $\theta_{13} = 0$, once triggered by subleading effects.

Another fundamental point is that SN neutrino oscillations are not sensitive to arbitrarily small values of the mixing angle. The fantastic sensitivity to an arbitrarily small mixing angle, as it appears in two-flavor analyses, disappears when one takes into account other sub-leading corrections. As a result, strategies outlined in Refs. [15, 16] may be useful for determination of the mass hierarchy if the relevant signals are observed, but not for determination of a non-zero θ_{13} itself. On the other hand, in principle we could determine the neutrino mass hierarchy even if θ_{13} were exactly zero—which

might end up being our only hope if θ_{13} is beyond the reach of laboratory-based oscillation experiments.

Acknowledgements

This work was partly supported by the Deutsche Forschungsgemeinschaft under grant TR-27 “Neutrinos and Beyond” and the Cluster of Excellence “Origin and Structure of the Universe” (Munich and Garching). The work of I.T. has been partly supported by the Italian MIUR and INFN through the “Astroparticle Physics” research project. Her stay in Munich has been partly supported by a junior fellowship awarded by the Italian Society of Physics (Borsa SIF “Antonio Stanghellini”).

-
- [1] J. T. Pantaleone, Phys. Lett. B **287**, 128 (1992).
 - [2] G. Sigl and G. Raffelt, Nucl. Phys. B **406**, 423 (1993).
 - [3] V. A. Kostelecký and S. Samuel, Phys. Rev. D **52**, 621 (1995).
 - [4] S. Pastor, G. G. Raffelt and D. V. Semikoz, Phys. Rev. D **65**, 053011 (2002).
 - [5] R. F. Sawyer, Phys. Rev. D **72**, 045003 (2005).
 - [6] H. Duan, G. M. Fuller and Y.-Z. Qian, Phys. Rev. D **74**, 123004 (2006).
 - [7] H. Duan, G. M. Fuller, J. Carlson and Y.-Z. Qian, Phys. Rev. D **74**, 105014 (2006).
 - [8] S. Hannestad, G. G. Raffelt, G. Sigl and Y. Y. Y. Wong, Phys. Rev. D **74**, 105010 (2006); Erratum *ibid.* **76**, 029901 (2007).
 - [9] H. Duan, G. M. Fuller, J. Carlson and Y.-Z. Qian, Phys. Rev. D **75**, 125005 (2007).
 - [10] G. G. Raffelt and A. Yu. Smirnov, Phys. Rev. D **76**, 081301 (2007); Erratum *ibid.* **77**, 029903 (2008).
 - [11] G. L. Fogli, E. Lisi, A. Marrone and A. Mirizzi, JCAP **0712**, 010 (2007).
 - [12] G. L. Fogli, E. Lisi, A. Marrone, A. Mirizzi and I. Tamborra, Phys. Rev. D **78**, 097301 (2008).
 - [13] B. Dasgupta, A. Dighe, G. G. Raffelt and A. Yu. Smirnov, Phys. Rev. Lett. **103**, 051105 (2009).
 - [14] G. Fogli, E. Lisi, A. Marrone and I. Tamborra, JCAP **0910** (2009) 002.
 - [15] H. Duan, G. M. Fuller, J. Carlson and Y.-Z. Qian, Phys. Rev. Lett. **99**, 241802 (2007).
 - [16] B. Dasgupta, A. Dighe and A. Mirizzi, Phys. Rev. Lett. **101**, 171801 (2008).
 - [17] H. Duan, G. M. Fuller, J. Carlson and Y.-Z. Qian, Phys. Rev. Lett. **100**, 021101 (2008).
 - [18] H. Duan, G. M. Fuller and Y.-Z. Qian, Phys. Rev. D **77**, 085016 (2008).
 - [19] B. Dasgupta, A. Dighe, A. Mirizzi and G. G. Raffelt, Phys. Rev. D **77**, 113007 (2008).
 - [20] A. Esteban-Pretel, A. Mirizzi, S. Pastor, R. Tomàs, G. G. Raffelt, P. D. Serpico and G. Sigl, Phys. Rev. D **78**, 085012 (2008).
 - [21] A. Friedland, arXiv:1001.0996 [hep-ph].
 - [22] A. Esteban-Pretel, PhD Thesis (University of Valencia, 2009), arXiv:0912.1616 [hep-ph].
 - [23] A. Esteban-Pretel, S. Pastor, R. Tomàs, G. G. Raffelt and G. Sigl, Phys. Rev. D **77**, 065024 (2008).
 - [24] B. Dasgupta and A. Dighe, Phys. Rev. D **77**, 113002 (2008).
 - [25] G. Fogli, E. Lisi, A. Marrone and I. Tamborra, JCAP **0904**, 030 (2009).
 - [26] J. Gava and C. Volpe, Phys. Rev. D **78**, 083007 (2008).
 - [27] M. Blennow, A. Mirizzi and P. D. Serpico, Phys. Rev. D **78**, 113004 (2008).
 - [28] F. J. Botella, C. S. Lim and W. J. Marciano, Phys. Rev. D **35**, 896 (1987).
 - [29] A. Mirizzi, S. Pozzorini, G. G. Raffelt and P. D. Serpico, JHEP **0910**, 020 (2009).
 - [30] R. C. Schirato and G. M. Fuller, arXiv:astro-ph/0205390.